

Phase transition of the higher dimensional charged Gauss-Bonnet black hole in de Sitter spacetime

Meng-Sen Ma^{a,b*}, Li-Chun Zhang^{a,b†}, Hui-Hua Zhao^{a,b‡}, Ren Zhao^{a,b§}

^a*Department of Physics, Shanxi Datong University, Datong 037009, China*

^b*Institute of Theoretical Physics, Shanxi Datong University, Datong 037009, China*

We study the phase transition of charged Gauss-Bonnet-de Sitter (GB-dS) black hole. For black holes in de Sitter spacetime, there is not only black hole horizon, but also the cosmological horizon. The thermodynamic quantities on the both horizons satisfy the first law of the black hole thermodynamics, respectively; moreover, there are additional connections between them. Using the effective temperature approach, we obtained the effective thermodynamic quantities of charged GB-dS black hole. According to Ehrenfest classification, we calculate some response functions and plot their figures, from which one can see that the spacetime undergoes a second-order phase transition at the critical point. It is shown that the critical values of effective temperature and pressure decrease with the increase of the value of GB parameter α .

PACS numbers: 04.70.Dy

I. INTRODUCTION

Due to the existence of Hawking radiation and the entropy, black holes may display themselves like thermodynamic systems[1–4]. Phase transitions and critical phenomena are important characteristics of ordinary thermodynamic system. Thus, the natural question to ask is whether there also exists phase transition in the black hole thermodynamics. The pioneer work of Hawking and Page give us an definitive answer[5]. Recently, the phase transitions of black holes in asymptotically anti de-Sitter (AdS) spacetime have received considerable attention[6–15]. In particular, by considering the cosmological constant as thermodynamic pressure $P = -\frac{\Lambda}{8\pi} = \frac{(d-1)(d-2)}{16\pi l^2}$, one can introduce an extended phase space, in which the $P - V$ criticality has been extensively discussed[16–29, 31, 36]. It is shown that many black holes in AdS spacetime exhibit similar critical behaviors to that of Van der Waals liquid-gas system.

The astronomical observations show that our Universe is probably an asymptotically de Sitter (dS) one. It raises the interest on black holes in dS spacetime. There are many works on the thermal properties of black holes in de Sitter(dS) spacetime[32–43]. As is well known, there are multi-horizons for black holes in dS spacetime. On the black hole event horizon and the cosmological horizon there are different temperatures, which prevent the black hole thermodynamic system in equilibrium. It should be noted that there are common parameters M , Q , Λ in the thermodynamic quantities corresponding to the black hole horizon and that corresponding to the cosmological horizon. Thus, these thermodynamic quantities are not independent. Taking into account their connections is relevant to the understanding of thermal properties of dS spacetime.

Higher derivative curvature terms occur in many occasions, such as in the semiclassical quantum gravity and in the effective low-energy action of superstring theories. Among the many theories of gravity with higher derivative curvature terms, due to the special features the Gauss-Bonnet (GB) gravity has attract much interest. The thermodynamic properties and phase structures of GB-AdS black hole have been briefly discussed in [44]. In Refs.[23–25], the critical phenomena and phase transition of the charged GB-AdS black hole have been studied extensively.

In this paper, we study the thermal properties of charged GB-dS black hole after considering the connections between the black hole horizon and the cosmological horizon. Using the effective equilibrium temperature approach, we calculate some effective thermodynamic quantities, from which we can analyze whether there exists phase transition for charged GB-dS black hole.

The paper is arranged as follows: in the next section we simply introduce the charged Gauss-Bonnet-dS black hole. In section 3, we will calculate the effective thermodynamic quantities. In section 4, we derive the heat capacity at constant pressure, the volume expansivity β , and the isothermal compressibility κ and plot some curves to demonstrate the phase transition. We will make some concluding remarks in section 5. (we use the units $G_d = \hbar = c = k_B = 1$)

* Email:mengsenma@gmail.com

† Email:zhao2969@sina.com

‡ Email:kietemap@126.com

§ Email:zhao2969@sina.com

II. CHARGED GAUSS-BONNET-DS BLACK HOLE

The action of the d - dimensional Einstein-Gauss-Bonnet-Maxwell-dS theory has the form

$$S = \frac{1}{16\pi} \int d^d x \sqrt{-g} [R - 2\Lambda + \alpha(R_{\mu\nu\gamma\delta}R^{\mu\nu\gamma\delta} - 4R_{\mu\nu}R^{\mu\nu} + R^2) - 4\pi F_{\mu\nu}F^{\mu\nu}], \quad (1)$$

where α is the Gauss-Bonnet coupling constant and the cosmological constant is $\Lambda = \frac{(d-1)(d-2)}{2l^2}$ for dS spacetime, $F_{\mu\nu}$ is the Maxwell field strength. The d -dimensional static charged Gauss-Bonnet-dS black hole solution for the action is described by

$$ds^2 = -f dt^2 + f^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2 + \cos^2\theta d\Omega_{d-4}^2), \quad (2)$$

with the metric function given by [23, 24, 29]

$$f(r) = 1 + \frac{r^2}{2\tilde{\alpha}} \left(1 - \sqrt{1 + \frac{64\pi\tilde{\alpha}M}{(d-2)\Sigma_k r^{d-1}} - \frac{2\tilde{\alpha}Q^2}{(d-2)(d-3)r^{2d-4}} + \frac{8\tilde{\alpha}\Lambda}{(d-1)(d-2)}} \right), \quad (3)$$

where $\tilde{\alpha} = (d-3)(d-4)\alpha$, Σ_k is the area of a unite $(d-2)$ -dimensional sphere, M is the black hole mass, Q is related to charge of the black hole. Horizons occur where $f(r) = 0$, the largest root is the cosmological horizon $r = r_c$, and the next root is the black hole event horizon $r = r_+$. When $M = Q = 0$, the solution (2) reduces to the pure GB-dS space with only one cosmological horizon at $r_c = l_c$.

The equations $f(r_+) = 0$ and $f(r_c) = 0$ are rearranged to

$$M = \frac{(d-2)\Sigma_k r_+^{d-3}}{16\pi} \left(1 + \frac{\tilde{\alpha}}{r_+^2} \right) - \frac{\Sigma_k r_+^{d-1}\Lambda}{8\pi(d-1)} + \frac{\Sigma_k Q^2}{32\pi(d-3)r_+^{d-3}},$$

$$M = \frac{(d-2)\Sigma_k r_c^{d-3}}{16\pi} \left(1 + \frac{\tilde{\alpha}}{r_c^2} \right) - \frac{\Sigma_k r_c^{d-1}\Lambda}{8\pi(d-1)} + \frac{\Sigma_k Q^2}{32\pi(d-3)r_c^{d-3}},$$

from which one can derive

$$M = \frac{(d-2)\Sigma_k r_+^{d-3} r_c^{d-3} (r_c^2 - r_+^2)}{16\pi(r_c^{d-1} - r_+^{d-1})} \left(1 + \frac{\tilde{\alpha}(r_c^2 + r_+^2)}{r_c^2 r_+^2} \right) + \frac{\Sigma_k Q^2 (r_c^{2d-4} - r_+^{2d-4})}{32\pi(d-3)r_c^{d-3} r_+^{d-3} (r_c^{d-1} - r_+^{d-1})}, \quad (4)$$

by eliminating the Λ , and

$$\Lambda = \frac{(d-1)(d-2)}{2(r_c^{d-1} - r_+^{d-1})} (r_c^{d-3} - r_+^{d-3} + \tilde{\alpha}(r_c^{d-5} - r_+^{d-5})) - \frac{(d-1)Q^2(r_c^{d-3} - r_+^{d-3})}{4(d-3)r_c^{d-3} r_+^{d-3} (r_c^{d-1} - r_+^{d-1})}$$

by eliminating M .

The surface gravities of black hole horizon and the cosmological horizon are

$$\begin{aligned} \kappa_+ = \frac{f'(r)}{2} \Big|_{r=r_+} &= -\frac{1}{r_+} + \frac{(d-1)r_c^{d-3}r_+(r_c^2 - r_+^2)}{2(r_+^2 + 2\tilde{\alpha})(r_c^{d-1} - r_+^{d-1})} \left(1 + \frac{\tilde{\alpha}(r_c^2 + r_+^2)}{r_c^2 r_+^2} \right) \\ &\quad - \frac{Q^2}{4(d-3)(r_+^2 + 2\tilde{\alpha})r_+^{2d-7}} \left(2 - \frac{(d-1)(r_c^{2d-4} - r_+^{2d-4})}{(d-2)r_c^{d-3}(r_c^{d-1} - r_+^{d-1})} \right), \end{aligned} \quad (5)$$

$$\begin{aligned} \kappa_c = \frac{f'(r)}{2} \Big|_{r=r_c} &= -\frac{1}{r_c} + \frac{(d-1)r_+^{d-3}r_c(r_c^2 - r_+^2)}{2(r_c^2 + 2\tilde{\alpha})(r_c^{d-1} - r_+^{d-1})} \left(1 + \frac{\tilde{\alpha}(r_c^2 + r_+^2)}{r_c^2 r_+^2} \right) \\ &\quad - \frac{Q^2}{4(d-3)(r_c^2 + 2\tilde{\alpha})r_c^{2d-7}} \left(2 - \frac{(d-1)(r_c^{2d-4} - r_+^{2d-4})}{(d-2)r_+^{d-3}(r_c^{d-1} - r_+^{d-1})} \right). \end{aligned} \quad (6)$$

The thermodynamic quantities for the two horizons satisfy the first law of black hole thermodynamic[33, 36]

$$\delta M = \frac{\kappa_+}{2\pi} \delta S_+ + \Phi_+ \delta Q + V_+ \delta P + \tilde{V}_+ d\tilde{\alpha}, \quad (7)$$

$$\delta M = \frac{\kappa_c}{2\pi} \delta S_c + \Phi_c \delta Q + V_c \delta P + \tilde{V}_c d\tilde{\alpha}, \quad (8)$$

where

$$\begin{aligned} S_+ &= \frac{\Sigma_k r_+^{(d-2)}}{4} \left(1 + \frac{2(d-2)\tilde{\alpha}}{(d-4)r_+^2} \right), \quad S_c = \frac{\Sigma_k r_c^{(d-2)}}{4} \left(1 + \frac{2(d-2)\tilde{\alpha}}{(d-4)r_c^2} \right), \\ \Phi_+ &= \frac{\Sigma_k Q}{16\pi(d-3)r_+^{d-3}}, \quad \Phi_c = \frac{\Sigma_k Q}{16\pi(d-3)r_c^{d-3}}, \quad V_+ = \frac{\Sigma_k r_+^{d-1}}{d-1}, \quad V_c = \frac{\Sigma_k r_c^{d-1}}{d-1}, \\ \tilde{V}_+ &= \frac{(d-2)}{16\pi} r_+^{d-5}, \quad \tilde{V}_c = \frac{(d-2)}{16\pi} r_c^{d-5}, \quad P = -\frac{\Lambda}{8\pi}. \end{aligned} \quad (9)$$

III. EFFECTIVE THERMODYNAMIC QUANTITIES OF CHARGED GAUSS-BONNET BLACK HOLE

We have given the thermodynamic quantities without considering the relationship between the black hole horizon and the cosmological horizon. Because there are four variables M , Q , Λ and $\tilde{\alpha}$ in the spacetime, the thermodynamic quantities corresponding to the black hole horizon and the cosmological horizon are functions of these variables. Through the four variables, there are some connections between the thermodynamic quantities corresponding to the black hole horizon and the ones corresponding to the cosmological horizon. When the thermodynamic property of charged de Sitter spacetime is studied, we must consider the relationship with the two horizon. Recently, by studying Hawking radiation of de Sitter spacetime, refs. [45–47] obtained that the outgoing rate of the charged de Sitter spacetime which radiates particles with energy ω is

$$\Gamma = e^{\Delta S_+ + \Delta S_c}, \quad (10)$$

where ΔS_+ and ΔS_c are Bekenstein-Hawking entropy difference corresponding to the black hole horizon and the cosmological horizon after the charged de Sitter spacetime radiates particles with energy ω . Therefore, the thermodynamic entropy of the charged de Sitter spacetime is the sum of the black hole horizon entropy and the cosmological horizon entropy:

$$S = S_+ + S_c. \quad (11)$$

Recently, thermodynamic volume of charged dS black hole is given by[33, 36]

$$V = V_c - V_+. \quad (12)$$

From (8), one can obtain

$$dS_+ = \frac{\Sigma_k(d-2)}{4} r_+^{d-5} (r_+^2 + 2\tilde{\alpha}) dr_+, \quad dS_c = \frac{\Sigma_k(d-2)}{4} r_c^{d-5} (r_c^2 + 2\tilde{\alpha}) dr_c. \quad (13)$$

Substituting (11) and (12) into (13), we obtain

$$\begin{aligned} dS_+ &= \frac{r_+^3(r_+^2 + 2\tilde{\alpha})dS}{r_c^3(r_+^2 + 2\tilde{\alpha}) + r_+^3(r_c^2 + 2\tilde{\alpha})} - \frac{(d-2)(r_+^2 + 2\tilde{\alpha})(r_c^2 + 2\tilde{\alpha})dV}{4[r_c^3(r_+^2 + 2\tilde{\alpha}) + r_+^3(r_c^2 + 2\tilde{\alpha})]}, \\ dS_c &= \frac{r_+^3(r_c^2 + 2\tilde{\alpha})dS}{r_c^3(r_+^2 + 2\tilde{\alpha}) + r_+^3(r_c^2 + 2\tilde{\alpha})} + \frac{(d-2)(r_+^2 + 2\tilde{\alpha})(r_c^2 + 2\tilde{\alpha})dV}{4[r_c^3(r_+^2 + 2\tilde{\alpha}) + r_+^3(r_c^2 + 2\tilde{\alpha})]}. \end{aligned} \quad (14)$$

Substituting (14) into (7) and (8), we get

$$dM = T_{eff}dS - P_{eff}dV + \Phi_{eff}dQ + \tilde{V}_{eff}d\tilde{\alpha}, \quad (15)$$

where the effective temperature T_{eff} , effective electric potential Φ_{eff} and effective pressure P_{eff} are respectively

$$\begin{aligned} T_{eff} &= \frac{\kappa_+ V_c r_c^3 (r_+^2 + 2\tilde{\alpha}) - \kappa_- V_+ r_+^3 (r_c^2 + 2\tilde{\alpha})}{2\pi(V_c - V_+)[r_c^3(r_+^2 + 2\tilde{\alpha}) + r_+^3(r_c^2 + 2\tilde{\alpha})]}, \quad \Phi_{eff} = \frac{\Phi_+ V_c - \Phi_- V_+}{V_c - V_+}, \\ P_{eff} &= \frac{(d-2)(r_+^2 + 2\tilde{\alpha})(r_c^2 + 2\tilde{\alpha})}{8\pi[r_c^3(r_+^2 + 2\tilde{\alpha}) + r_+^3(r_c^2 + 2\tilde{\alpha})]} \frac{(\kappa_+ V_c + \kappa_- V_+)}{(V_c - V_+)}. \\ \tilde{V}_{eff} &= \frac{V_c \tilde{V}_+ - V_+ \tilde{V}_c}{V_c - V_+} \end{aligned} \quad (16)$$

Substituting (5), (6) and (9) into (16), we obtain

$$\begin{aligned} T_{eff} &= \frac{x[(d-1)(1-x^2)(1+x^{d-1}) - 2(1-x^{d+1})]}{4\pi r_c(1-x^{d-1})[(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)]} \\ &+ \frac{\tilde{\alpha}[(d-1)(1-x^2)(1+x^{d-1})(1+x^2) - 4(1-x^{d+3})]}{4\pi r_c^2 x(1-x^{d-1})[(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)]} \\ &+ \frac{Q^2}{8\pi(d-3)(d-2)} \frac{(d-1)(1+x^{d-1})(1-x^{2d-4}) - 2(d-2)(1-x^{3d-5})}{r_c^{2d-5} x^{2d-7}(1-x^{d-1})[(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)]}, \end{aligned} \quad (17)$$

$$\begin{aligned} P_{eff} &= -\frac{(d-2)(x^2 + 2\tilde{\alpha}/r_c^2)(1 + 2\tilde{\alpha}/r_c^2)(1+x^d)}{4\pi r_c^2 x(1-x^{d-1})[(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)]} \\ &+ \frac{(d-2)(d-1)(1-x^2)x}{8\pi r_c^2(1-x^{d-1})^2} \left(\frac{(1+x^{2d-3}) + 2\tilde{\alpha}(1+x^{2d-5})/r_c^2}{(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)} \right) \\ &+ \frac{\tilde{\alpha}(d-2)(d-1)(1-x^2)(1+x^2)}{8\pi r_c^4(1-x^{d-1})^2 x} \left(\frac{(1+x^{2d-3}) + 2\tilde{\alpha}(1+x^{2d-5})/r_c^2}{(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)} \right) \\ &- \frac{Q^2(d-2)}{8\pi(d-3)r_c^{2d-4}x^{2d-7}(1-x^{d-1})} \left(\frac{(1+x^{3d-6}) + 2\tilde{\alpha}(1+x^{3d-8})/r_c^2}{(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)} \right) \\ &+ \frac{Q^2(d-1)(1-x^{2d-4})}{16\pi(d-3)r_c^{2d-4}x^{2d-7}(1-x^{d-1})^2} \left(\frac{(1+x^{2d-3}) + 2\tilde{\alpha}(1+x^{2d-5})/r_c^2}{(x^2 + 2\tilde{\alpha}/r_c^2) + x^3(1 + 2\tilde{\alpha}/r_c^2)} \right), \end{aligned} \quad (18)$$

$$\Phi_{eff} = \frac{\Sigma_k Q(1-x^{2d-4})}{16\pi(d-3)r_c^{d-3}x^{d-3}(1-x^{d-1})}. \quad (19)$$

where $x := r_+/r_c$ and $0 < x < 1$.

In the case of neglecting the connections between the two horizons, the black hole horizon and the cosmological horizon are two independent thermodynamic systems. Due to different horizon temperatures, the spacetime cannot be in thermal equilibrium. After considering the connections between the both horizons, from Eq.(17) one can see that only one effective temperature T_{eff} left.

IV. PHASE TRANSITION IN GAUSS-BONNET-DS BLACK HOLE SPACETIME

Recently, there are many works on the phase transition of black hole thermodynamic system. One can also compare the critical behaviors of van der Waals liquid-gas system and black hole system. Nevertheless, in dS spacetime there are the cosmological horizon and the black hole horizon. Generally, equilibrium cannot be achieved unless for extremal black hole. In this section, we employ the effective thermodynamic quantities to study the phase transition and critical behaviors of charged Gauss-Bonnet-dS black hole. This may avoid the apprehension of equilibrium of the black hole system.

Comparing with the Van der Waals equation

$$\left(P + \frac{a}{v^2}\right)(v - \tilde{b}) = kT, \quad (20)$$

Here, $v = V/N$ is the specific volume of the fluid, P its pressure, T its temperature, and k is the Boltzmann constant. Similarly, for the GB-dS black hole we can set the specific volume as

$$v = r_c(1 - x). \quad (21)$$

According to Eq.(18), one can employ the two equations

$$\left(\frac{\partial P_{eff}}{\partial x}\right)_{T_{eff}} = 0, \quad \left(\frac{\partial^2 P_{eff}}{\partial x^2}\right)_{T_{eff}} = 0, \quad (22)$$

to calculate the critical temperature, critical pressure and the critical specific volume. We can ascertain the critical values (r_c^0, x^0) for given Q, n and $\tilde{\alpha}$. According to these values, we can obtain the critical temperature, critical electric potential, critical pressure and critical volume. As is shown Table I, the critical temperature and critical pressure increase with the dimension of spacetime for fixed Q and $\tilde{\alpha}$. However, they decrease with the increase of the Gauss-Bonnet parameter $\tilde{\alpha}$ for fixed spacetime dimension and electric charge Q .

	T_{eff}^c			P_{eff}^c		
	$n = 5$	$n = 6$	$n = 7$	$n = 5$	$n = 6$	$n = 7$
$\tilde{\alpha} = 0.1$	0.028628	0.040562	0.049202	0.008229	0.018877	0.031920
$\tilde{\alpha} = 0.01$	0.033651	0.045736	0.054251	0.010356	0.021471	0.034484
$\tilde{\alpha} = 0.001$	0.034297	0.046385	0.054889	0.010632	0.021802	0.034836

TABLE I: Numerical solutions for T_{eff}^c , P_{eff}^c with fixed $Q = 1$ and spacetime dimension $n = 5, 6, 7$ and GB parameter $\tilde{\alpha} = 0.1, 0.01, 0.001$, respectively.

In order to describe the relation of P_{eff} and V in the vicinity of critical temperature, we plot the curves of P_{eff} - V at different temperatures. When the effective temperature $T_{eff} > T_{eff}^c$, the stable condition $(\frac{\partial P}{\partial V})_{T_{eff}} < 0$ can be satisfied. In the case of $T_{eff} < T_{eff}^c$, there exist a portion in the curve where $(\frac{\partial P}{\partial V})_{T_{eff}} > 0$, thus in these parts the system is unstable. So phase transition may occur only at $T_{eff} = T_{eff}^c$.

According to Ehrenfest classification, when the chemical potential and its first derivative are continuous, whereas the second derivative of chemical potential is discontinuous, this kind of phase transition is called the second-order phase transition. For Van der Waals equation, at the critical point no latent and sudden change in volume happen between liquid phase and gas phase. According to Ehrenfest classification, this kind of phase transition is continuous.

To understand the types of the phase transition for the GB-dS black hole, we should calculate the specific heat at constant pressure C_P , the volume expansivity β , and the isothermal compressibility κ :

$$C_P = T_{eff} \left(\frac{\partial S}{\partial T_{eff}} \right)_{P_{eff}} = T_{eff} \left(\frac{\partial S}{\partial x} \frac{\partial x}{\partial T_{eff}} \right)_{P_{eff}}, \quad (23)$$

$$\beta = \frac{1}{v} \left(\frac{\partial v}{\partial T_{eff}} \right)_{P_{eff}} = \frac{1}{v} \left(\frac{\partial v}{\partial x} \frac{\partial x}{\partial T_{eff}} \right)_{P_{eff}}, \quad (24)$$

$$\kappa_T = -\frac{1}{v} \left(\frac{\partial v}{\partial P_{eff}} \right)_{T_{eff}} = -\frac{1}{v} \left(\frac{\partial v}{\partial x} \frac{\partial x}{\partial P_{eff}} \right)_{T_{eff}}. \quad (25)$$




FIG. 1: $P_{eff} - V$ diagram of charged GB-dS black hole. From top to bottom the curves correspond to the effective temperature $T_{eff}^c + 0.0005$, $T_{eff}^c + 0.0002$, T_{eff}^c , $T_{eff}^c - 0.0002$ and $T_{eff}^c - 0.0005$. We have set $Q = 1$.

We also depict the curves of $C_P - x$, $\kappa_T - x$ and $\beta - x$ in the Fig.2, Fig.3 and Fig.4 respectively. From these curves, we find that, for the charged GB-dS black hole, there exists an infinite peak in these curves. Moreover, in Fig.5, one can see that the curves of $S - T_{eff}$ are continuous at the critical point. According to Ehrenfest, the phase transition of the charged GB-dS black hole should be the second-order one, which is similar to the RN-(A)dS black hole.




FIG. 2: $C_P - x$ curves for charged GB-dS black hole with $Q = 1$.

V. DISCUSSION AND CONCLUSION

In this paper, we adopted Ehrenfests classification to study the phase transition of the charged GB-dS black hole. After introducing the connection between the thermodynamic quantities corresponding to the black hole horizon and the cosmological horizon, we give the effective thermodynamic quantities, such as effective pressure P_{eff} , effective temperature T_{eff} , etc, of the black hole. From the equation of state described by the effective thermodynamic quantities, we study the $P - V$ criticality. As illustrated in Fig.1, it exhibits a similar phase transition to Van der Waals equation.

By treating the effective cosmological constant as the thermodynamic pressure, in the extended phase space, we completely follow the Ehrenfest classification to explore the type of the phase transition of the charged GB-dS black




FIG. 3: $\kappa_T - x$ curves for charged GB-dS black hole with $Q = 1$.




FIG. 4: $\beta - x$ curves for charged GB-dS black hole with $Q = 1$.

hole. It is shown that when the entropy is a continuous function of temperature. However, the heat capacity C_P , the isothermal compressibility κ_T and the expansion coefficient β are all divergent at the critical point. This means this kind of phase transition for the charged GB-dS black hole is second order.




FIG. 5: $S - T_{eff}$ with $Q = 1$.

Acknowledgments

This work is supported by NSFC under Grant Nos.(11475108,11175109;11075098;11205097) and the doctoral Sustentation foundation of Shanxi Datong University (2011-B-03).

-
- [1] J. D. Bekenstein, “Black holes and entropy”, *Phys. Rev. D* 7, 949 (1973).
 - [2] J. M. Bardeen, B. Carter, S. W. Hawking, “The Four laws of black hole mechanics”, *Commun. Math. Phys.* 31, 161 (1973).
 - [3] S. W. Hawking, “Black Hole Explosions”, *Nature*, 248 30 (1974).
 - [4] S. W. Hawking, “Particle Creation by Black Holes”, *Commun. Math. Phys.* 43,199(1975).
 - [5] S. Hawking and D. N. Page, “Thermodynamics of black holes in anti-de Sitter space”, *Commun. Math. Phys.* 87, 577 (1983).
 - [6] A. Chamblin, R. Emparan, C. Johnson, and R. Myers, “Charged AdS black holes and catastrophic holography”, *Phys.Rev. D*60 (1999) 064018, [hep-th/9902170].
 - [7] A. Chamblin, R. Emparan, C. Johnson, and R. Myers, “Holography, thermodynamics and uctuations of charged AdS black holes”, *Phys.Rev. D*60 (1999) 104026,[hep-th/9904197].
 - [8] A. Sahay, T. Sarkar, and G. Sengupta, “Thermodynamic geometry and phase transitions in Kerr-Newman-AdS black holes”, *JHEP*1004, 118 (2010), [arXiv:1002.2538[hep-th]].
 - [9] A. Sahay, T. Sarkar, and G. Sengupta, “On the thermodynamic geometry and critical phenomena of AdS black holes”, *JHEP*1007, 082 (2010), [arXiv:1004.1625[hep-th]].
 - [10] A. Sahay, T. Sarkar, and G. Sengupta, “On the phase structure and thermodynamic geometry of R-charged black holes”, *JHEP*1011, 125, (2010), [arXiv:1009.2236[hep-th]].
 - [11] R. Banerjee, S. Ghosh, and D. Roychowdhury, “New type of phase transition in Reissner Nordstr?mCAdS black hole and its thermodynamic geometry”, *Phys. Lett. B* 696, 156 (2011), [arXiv:1008.2644[gr-qc]].
 - [12] R. Banerjee, S. K. Modak, S. Samanta, “Second order phase transition and thermodynamic geometry in Kerr-AdS black holes”, *Phys. Rev. D*84, 064024 (2011), [arXiv:1005.4832[hep-th]].
 - [13] R. Banerjee, D. Roychowdhury, “Thermodynamics of phase transition in higher dimensional AdS black holes”, *JHEP*11, 004 (2011), [arXiv:1109.2433[hep-th]].
 - [14] R. Banerjee, D. Roychowdhury, “Critical phenomena in Born-Infeld AdS black holes”, *Phys. Rev. D* 85, 044040 (2012), [arXiv:1111.0147[hep-th]].
 - [15] R. Banerjee, D. Roychowdhury, “Critical behavior of Born-Infeld AdS black holes in higher dimensions”, *Phys. Rev. D* 85, 104043 (2012), [arXiv:1203.0118[hep-th]].
 - [16] D. Kastor, S. Ray, and J. Traschen, “Enthalpy and the mechanics of AdS black holes”, *Class. Quant. Grav.* 26, 195011 (2009), [arXiv:0904.2765[hep-th]].
 - [17] B. P. Dolan, “The cosmological constant and the black hole equation of state”, *Class.Quant.Grav.*28:125020,2011.
 - [18] S. Gunasekaran, D. Kubiznak, R. B. Mann, “Extended phase space thermodynamics for charged and rotating black holes and Born-Infeld vacuum polarization”, *JHEP*, 2012(11), 1-43, arXiv:1208.6251
 - [19] B. P. Dolan, “Compressibility of rotating black holes”, *Phys. Rev. D*84 (2011) 127503;
 - [20] M. Cvetič, G.W. Gibbons, D. Kubiznak, C.N. Pope, “Black Hole Enthalpy and an Entropy Inequality for the Thermodynamic Volume”, *Phys.Rev.D* 84:024037,2011, arXiv:1012.2888.
 - [21] David Kubiznak, Robert B. Mann, “P-V criticality of charged AdS black holes”, *JHEP* 1207:033,2012, arXiv:1205.0559.
 - [22] B. P. Dolan, “Where is the PdV term in the first law of black hole thermodynamics?”, [arXiv:1209.1272[gr-qc]].
 - [23] Shao-Wen Wei, Yu-Xiao Liu, “Critical phenomena and thermodynamic geometry of charged Gauss-Bonnet AdS black holes”, *Phys. Rev. D* 87, 044014 (2013), arXiv:1209.1707.
 - [24] Rong-Gen Cai, Li-Min Cao, Li Li, and Run-Qiu Yang, “P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space”, *JHEP*,2013(9), 1-22. [arXiv:1306.6233[gr-qc]].
 - [25] Wei Xu, Hao Xu, Liu Zhao, “Gauss-Bonnet coupling constant as a free thermodynamical variable and the associated criticality”, *Eur. Phys. J. C* (2014) 74:2970.
 - [26] S. H. Hendi and M. H. Vahidinia, “Extended phase space thermodynamics and $P?V$ criticality of black holes with a nonlinear source”, *Phys. Rev. D*88,084045(2013), arXiv:1212.6128[hep-th];
 - [27] R. Zhao, H. H. Zhao, M. S. Ma and L. C. Zhang, “On the critical phenomena and thermodynamics of charged topological dilaton AdS black holes”, *Eur.Phys. J. C.*,2013, 73:2645;
 - [28] N. Altamirano, D. Kubiznak and R. Mann, “Reentrant phase transitions in rotating antiCde Sitter black holes”, *Phys. Rev. D*,88, 101502(2013),arXiv:1306.5756 [hep-th].
 - [29] S. W. Wen, Y. X. Liu, “Triple points and phase diagrams in the extended phase space of charged Gauss-Bonnet black holes in AdS space”, *Phys. Rev. D*90, 044057 (2014).
 - [30] B. P. Dolan, “Vacuum energy and the latent heat of AdS-Kerr black holes”, *Phys. Rev. D*90, 084002 (2014).
 - [31] B. Mirza, Z. Sherkatghanad, “Phase transitions of hairy black holes in massive gravity and thermodynamic behavior of charged AdS black holes in an extended phase space”, *Phys. Rev. D*90, 084006 (2014).
 - [32] R.G. Cai, “Cardy-Verlinde formula and thermodynamics of black holes in de Sitter spaces”, *Nucl. Phys. B*628 (2002)375
 - [33] Y. Sekiwa, “Thermodynamics of de Sitter black holes: Thermal cosmological constant”, *Phys. Rev. D* 73(2006)084009; hep-th/0602269.

- [34] M. Urano and A. Tomimatsu, “Mechanical First Law of Black Hole Spacetimes with Cosmological Constant and Its Application to Schwarzschild-de Sitter Spacetime”, *Class.Quant.Grav.* 26(2009)105010, arXiv:0903.4230.
- [35] L. C. Zhang, H. F. Li and R. Zhao, “ Thermodynamics of the Reissner-Nordström-de Sitter black hole”, *Science Chian, Physics, Mechanics Astronomy*, 54(2011) 1384.
- [36] B. P. Dolan, D. Kastor, D. Kubiznak, R. B. Mann, J. Traschen, “Thermodynamic Volumes and Isoperimetric Inequalities for de Sitter Black Holes”, *Phys. Rev. D*, 87,104017(2013), arXiv:1301.5926.
- [37] Yun Soo Myung, “ Thermodynamics of the Schwarzschild-de Sitter black hole: Thermal stability of the Nariai black hole”, *Phys. Rev. D* 77(2008) 104007
- [38] R. G. Cai, *Physics* 34(2005) 555(in Chinese)
- [39] S. Bhattacharya, A. Lahiri, “Mass function and particle creation in Schwarzschild-de Sitter spacetime”, *The European Physical Journal C*, 2013, 73(12): 1-10, arXiv:1301.4532.
- [40] Hui-Hua Zhao, Li-Chun Zhang, Meng-Sen Ma, Ren Zhao, “ $P - V$ criticality of higher dimensional charged topological dilaton de Sitter black holes”, *Phys. Rev. D* 90, 064018.
- [41] R.G. Cai, J. Y. Ji, K. S. Soh, “ Action and entropy of black holes in spacetimes with a cosmological constant”, *Classical. Quantum. Grav.* 15 (1998) 2783.
- [42] G. W. Gibbons, H. Lü, D. N. Page, and C. N. Pope, “Rotating black holes in higher dimensions with a cosmological constant”, *Phys. Rev. Lett.* 93 (2004) 171102, [hep-th/0409155].
- [43] G. W. Gibbons, H. Lü, D. N. Page, and C. N. Pope, “The general Kerr-de Sitter metrics in all dimensions”, *J.Geom. Phys.* 53 (2005) 49–73, [hep-th/0404008]
- [44] Rong-Gen Cai, “ Gauss-Bonnet black holes in AdS”, *Phys. Rev. D* 65,084014(2002).
- [45] R. Zhao, L. C. Zhang, H. F. Li. “Hawking radiation of Reisser-Nordstrom-de Sitter space”, *Gen. Relat. Grav.* 42(2010)975
- [46] R. Zhao, L. C. Zhang, H. F. Li, Y.Q. Wu, “ Hawking radiation of a high-dimensional rotating black hole ”, *Eur. Phys. J. C* 65, 289(2010). (2010).
- [47] R. Zhao, L. C. Zhang, “ A new explanation for statistical entropy of charged black hole”, *Sci China-Phys Mech Astron*, 2013, 56(9): 1632-1635.